

INDEX TO VOLUME 58

AUTHOR

- Alton, Miles S. and Christine J. Blackburn: Diel changes in the vertical distribution of the euphausiids, *Thysanoessa spinifera* Holmes and *Euphausia pacifica* Hansen, in coastal waters of Washington, 179-190
- Azevedo, John A., Jr., Eldredge G. Hunt and Leon A. Woods, Jr.: Melanistic mutant in ringneck pheasants, 175-178
- Barlow, George W., and Victor L. De Vlaming: Ovarian cycling in long-jaw gobies, *Gillichthys mirabilis*, from the Salton Sea, 50-57
- Barnhart, Roger A.: see Kesner and Barnhart, 204-220
- Benville, Pete E., Jr.: see Earnest and Benville, 127-132
- Blackburn, Christine J.: see Alton and Blackburn, 179-190
- Borneman, John C.: see Carrier, Mallette, Wilbur and Borneman, 327-328; see Mallette, Wilbur, Carrier and Borneman, 67-68
- Briggs, Kenneth T., and C. William Davis: A study of predation by sea lions on salmon in Monterey Bay, 37-43
- Bury, C. Bruce: The effects of diesel fuel on a stream fauna, 291-295
- Cappucci, D. T., Jr. and W. M. Longhurst: Rabies in deer, 141-144
- Carrier, W. Dean: see Mallette, Wilbur, Carrier and Borneman, 67-68
- Carrier, W. Dean, Robert D. Mallette, Sanford Wilbur and John C. Borneman: California condor survey, 327-328
- Castle, William T. and Leon A. Woods, Jr.: DDT residues in white croakers, 198-203
- Chamberlain, Lawrence L.: Primary productivity in a new and older California reservoir, 254-267; see Lasker, Tenaza and Chamberlain, 58-66
- Chen, Lo-Chai: see Rosenblatt and Chen, 32-36
- Clark, William E.: see Espinosa and Clark, 149-152
- Davis, C. William: see Briggs and Davis, 37-43
- Day, John S.: see St. Amant and Day, 154-155
- Demory, Robert L.: Tailless Dover sole from off the Oregon coast, 147-148
- De Vlaming, Victor L.: see Barlow and De Vlaming, 50-57
- Dexter, Deborah M.: Molting and growth in laboratory reared phyllosomes of the California spiny lobster, *Panulirus interruptus*, 107-115
- Duffy, John M.: see Houk and Duffy, 321-323
- Earnest, Russell D. and Pete E. Benville, Jr.: Acute toxicity of four organochlorine insecticides to two species of surf perch, 127-132
- Elliott, George V. and T. M. Jenkins, Jr.: Winter food of trout in three high elevation Sierra Nevada lakes, 231-237
- Espinosa, Lawrence R. and William E. Clark: A polypropylene light trap for aquatic invertebrates, 149-152
- Farley, David G.: A range extension for the logperch, 248
- Fitch, John E.: A case for striped mullet, *Mugil cephalus*, spawning at sea, 246-248; The cottonmouth jack, *Uraspis secunda*, added to the marine fauna of California, 245-246
- Franklin, George W.: see Schneegas and Franklin, 133-140
- Grover, Charles A.: Population differences in the swell shark *Cephaloscyllium ventriosum*, 191-197
- Haaker, Peter L.: First record of a reversed butter sole, *Isopsetta isolepis*, 244-245
- Hanson, Jack A.: Tolerance of high salinity by the pileworm, *Neanthes succinea*, from the Salton Sea, California, 152-154
- Hashagen, Kenneth A., Jr.: see Rawstron and Hashagen, 221-230
- Hawthorne, Vernon M.: Coyote food habits in Sagehen Creek basin, northeastern California, 5-12
- Hobson, Edmund S.: The survival of Guadalupe cardinalfish *Apogon guadalupensis* at San Clemente Island, 68-69
- Houk, James L. and John M. Duffy: Two new sea urchin-acorn barnacle associations, 321-323
- Hunt, Eldredge G.: see Azevedo, Hunt and Woods, 175-178

- Iverson, Ernest W.: New hosts and bathymetric range extension for *Colobomatus embiotocae* (Crustacea, Coppepoda), 323-325
- Jenkins, T. M., Jr.: see Elliott and Jenkins, 231-237
- Jones, Albert C.: Contributions to the life history of the Piute sculpin in Sagehen Creek, California, 285-290
- Kesner, William D. and Roger A. Barnhart: Characteristics of the fall-run steelhead trout (*Salmo gairdneri gairdneri*) of the Klamath River system with emphasis on the half-pounder, 204-220
- Knaggs, Eric H.: see Parrish and Knaggs, 13-21; Southern California Pacific mackerel fishery and age composition of commercial landings during the 1968-69 and 1969-70 seasons, 116-120
- Kukowski, Gary E.: Southern range extension for the yellowfin goby, *Acanthogobius flavimanus* (Temminck and Schlegel), 326-327
- Lasker, Reuben, Richard H. Tenaza, and Lawrence L. Chamberlain: The response of Salton Sea fish eggs and larvae to salinity stress, 58-66
- Lea, Robert N.: Southern geographical records for four surfperches, family Embiotocidae, with notes on a population resurgence of the sharpnose seaperch, 27-31
- Longhurst, W. M.: see Cappucci and Longhurst, 141-144
- Mais, Kenneth F.: A subpopulation study of the Pacific sardine, 296-314
- Mallette, Robert D.: see Carrier, Mallette, Wilbur and Borneman, 327-328
- Mallette, Robert D., Sanford Wilbur, W. Dean Carrier and John C. Borneman: California condor survey, 1970, 67-68
- Martin, Michael: Morphology and variation of the Modoc sucker, *Catostomus microps* Rutter, with notes on feeding adaptations, 277-284
- Miller, Lee W.: Migrations of sturgeon tagged in the Sacramento-San Joaquin estuary, 102-106; White sturgeon population characteristics in the Sacramento-San Joaquin estuary as measured by tagging, 94-101
- Moring, John R.: Check list of intertidal fishes of Trinidad Bay, California, and adjacent areas, 315-320
- Morrell, Stephen: Life history of the San Joaquin kit fox, 162-174
- Odenweller, Dan Bowman: A new range record for the umbrella crab, *Cryptolithodes sitchensis* Brandt, 240-243
- Parrish, Richard H.: Symbiosis in the blacktail snailfish, *Careproctus melanurus*, and the box crab, *Lopholithodes foraminatus*, 239-240
- Parrish, Richard H., and Eric H. Knaggs: The southern California Pacific mackerel fishery and age composition of the catch for the 1964-65 through 1967-68 seasons, 13-21
- Penhale, Leonard B.: Reproductive failure of pelagic cormorant, San Luis Obispo County, California, 1970, 238
- Rawstron, Robert R.: Harvest, survival, and cost of two domestic strains of tagged rainbow trout stocked in Lake Berryessa, California, 44-49; Nonreporting of tagged largemouth bass, 1966-1969, 145-147.
- Rawstron, Robert R. and Kenneth A. Hashagen, Jr.: Mortality and survival rates of tagged largemouth bass (*Micropterus salmoides*) at Merle Collins Reservoir, 221-230
- Rosenblatt, Richard H., and Lo-Chai Chen: The identity of *Sebastes babcocki* and *Sebastes rubrivinctus*, 32-36
- Schneeegas, Edward R. and George W. Franklin: The Mineral King deer herd, 133-140
- Shaw, Stanton B.: DDT residues in eight California marine fishes, 22-26
- Spratt, Jerome D.: Age and length composition of northern anchovies, *Engraulis mordax*, in the California anchovy reduction fishery for the 1969-70 season, 121-126
- St. Amant, James A. and John S. Day: Range extension of *Palaeomonetes paludosus* (Gibbes) in California, 154-155
- Tenaza, Richard H.: see Lasker, Tenaza and Chamberlain, 58-66
- Varoujean, Daniel H.: The reoccurrence of the California scorpionfish, *Scorpaena guttata* Girard, in Monterey Bay, 238-239
- von Geldern, C. E., Jr.: A midwater trawl for threadfin shad, *Dorosoma petenense*, 268-276; Angling quality at Folsom Lake, California, as determined by a roving creel census, 75-93
- Wilbur, Sanford: see Carrier, Mallette, Wilbur and Borneman, 327-328; see Mallette, Wilbur, Carrier and Borneman, 67-68
- Woods, Leon A., Jr.: see Azevedo, Hunt and Woods, 175-178; see Castle and Woods, 198-203

SCIENTIFIC NAMES

- Acanthogobius flavimanus*: 326-327
Accipiter cooperii: 67
 striatus: 67
Acipenser medirostris: 102-106
 transmontanus: 94-101, 102-106
Alces alces: 141
Amphistichus argenteus: 27
Anisotremus davidsoni: 58-66
Anoplopoma fimbria: 22-26
Apogon guadalupensis: 68-69
 retrosella: 68
Aquila chrysaetos: 67
Arctostaphylos sp.: 137
Artemia sp.: 107-115
Astacus sp.: 292
Bairdiella chrysura: 65
 icistia: 58-66
Balanus concavus pacificus: 321-322
 nubilis: 321-322
Buteo jamaicensis: 67
Cadulus fusiformis: 323
Canis latrans: 5-12
Capreolus capreolus: 141
Careproctus melanurus: 239-240
Cathartes aura: 67
Catostomus microps: 277-284
 platyrhynchus: 287
 rimiculus: 292
 tahoensis: 287
Cephaloscyllium ventriosum: 191-197
Cervus canadensis: 141
 elaphus: 141
Chaenogobius urotaenia: 54
Chara, sp.: 154
Chasmichthys dolichognathus: 54
Chromis punctipinnis: 30
Circus cyaneus: 67
Citellus beecheyi: 5-12
 beldingi: 5-12
 lateralis: 5-12
Citharichthys sordidus: 22-26
Cladophora sp.: 294
Clemmys marmorata: 294
Clostridium botulinum: 149
Colobomatus embiotocae: 323-325
Coryphaenoides acrolepis: 22-26
Cottus beldingi: 285-290
Crago sp.: 94
Cryptolithodes sitchensis: 240-243
Cymatogaster aggregata: 28, 127-132
Cynoscion xanthulus: 58-66, 152
Dama dama: 141
Dentalium pretiosum: 323
Dorosoma petenense: 44, 92, 268-276
Echidnophaga gallinacea: 170
Engraulis mordax: 121-126
Entosphenus tridentatus: 292
Eopsetta jordani: 22-26
Erethizon dorsatum: 5-12
Euarctos americanus: 137
Eumetopias jubata: 38
Euphausia pacifica: 179-190
Eutamias sp.: 5-12
Falco mexicanus: 67
 sparverius: 67
Fasciola hepatica: 142
Fundulus confluentus: 55
Genyonemus lineatus: 22-26, 198-203
Gillichthys mirabilis: 50-57
Glaucomyx sabrinus: 5-12
Gobius paganelius: 51
Gymnogyps californianus: 67-68, 327-328
Hippampus zosterae: 55
Hyperprosopon anale: 323
 ellipticum: 27
Ictalurus catus: 78-93
 melas: 78-93
 nebulosus: 78-93
Isopsetta isolepis: 244-245
Lasaea cistula: 324
Lepomis cyanellus: 78-93
 macrochirus: 75-93
 microlophus: 78-93
Lepus americanus: 5-12
 sp.: 5-12
Lipoptena depressa: 142
Lopholithodes foraminatus: 239-240
Lytechinus sp.: 109
Macrocystis sp.: 68
Marmota flaviventris: 5-12, 136
Medialuna californiensis: 30
Mephitis mephitis: 143
Mergus merganser: 294
Micrometrus minimus: 127-132
Micropterus dolomieu: 75-93
 salmoides: 75-93, 145-147, 221-230
Microstomus pacificus: 147-148
Microtus sp.: 5-12
Mugil cephalus: 246-248
Mytilus sp.: 107-115
Nassarius mendicus: 323
Neanthes succinea: 152-154
Neolipoptena ferrisi: 142
Nuculana taphria: 323
Odocoileus hemionus: 141
 californicus: 133-140
 columbianus: 141-143
 virginianus: 141
Olivella sp.: 323
Oncorhynchus nerka: 87
 sp.: 37-43
Ophiodon elongatus: 22-26
Palaemonetes paludosus: 154-155
Panulirus interruptus: 107-115
Parophrys vetulus: 22-26
Percina caprodes: 248
Peromyscus sp.: 5-12
Phalacrocorax pelagicus: 238
Phanerodon atripes: 28-31
 furcatus: 30
Phasianus colchicus torquatus: 175-178

Phoradendron villosum: 136
Pocilia latipinna: 154
Pomoxis annularis: 86
nigromaculatus: 86
Prosopium williamsoni: 287
Pterogobius elapoides: 54
Pulex simulans: 142
Rana boylei: 294
Rangifer tarandus: 141
Rhacochilus toxotes: 28
vacca: 30
Rhinichthys osculus: 287, 292
Richardsonius egregius: 287
Salmo clarki henshawi: 44
gairdneri: 44-49, 78-93, 231-237, 285, 292
g. gairdneri: 204-220
trutta: 285
Salvelinus fontinalis: 231-237, 285
Sardinops caeruleus: 296-314
Scomber japonicus: 13-21, 116-120
Scorpaena guttata: 238-239

Scyllarus americanus: 107
Sebastes babcocki: 32-36
rubrivinctus: 32-36
Spirogyra sp.: 294
Strongylocentrotus franciscanus: 321-322
purpuratus: 321-322
Sylvilagus sp.: 5-12
Tamiasciurus douglasi: 5-12
Thamnophis couchi: 294
Thelazia californiensis: 142
Thysanoessa spinifera: 179-190
Tilapia mossambica: 154
Trachurus symmetricus: 13, 22-26, 116
Tubifex sp.: 109
Turbonilla sp.: 323
Uraspis secundus: 245-246
Urocyon cinereoargenteus: 143
Vulpes macrotis mutica: 162-174
Zalambius rosaceus: 323
Zalophus californianus: 37-43
Zygnuma sp.: 294

SUBJECT

Age: of anchovies taken the 1969-70 season, 121-126; of Klamath River fall-run steelhead, 204-220; of Pacific mackerel during 1964-65 through 1967-68 seasons, 13-21; of Pacific mackerel in southern California fishery, 1968-69, 1969-70, 116-120
Aldrin: toxicity to surfperch, 127-132
Anchovy, northern: fishery during 1969-70 season, 121-126
Angling: quality at Folsom Lake, 75-93
Bairdiella: effect of salinity on eggs and larvae, 58-66
Barnacle, acorn: association with sea urchins, 321-323
Bass: fishery at Folsom Lake, 83
Bass, largemouth: mortality and survival at Merle Collins Reservoir, 221-230; nonreporting of tagged fish, 145-147
Birds: as food for coyotes, 9
Cardinalfish, Guadalupe: survival at San Clemente Island, 68-69
Catfish: fishery at Folsom Lake, 84-85
Cattle: as food for coyotes, 7
Census, roving creel: of fishery at Folsom Lake, 75-93
Check list: intertidal fishes of Trinidad Bay and vicinity, 315-320
Chlorinated hydrocarbons: residues in California marine fishes, 22-26
Coleman Kamloops trout strain: performance at Lake Berryessa, 44-49
Condor, California: 1970 population survey, 67-68; population survey of 1971, 327-328
Copepod, parasitic: on embiotocid fish, 323-325

Cormorant, pelagic: reproductive failure in San Luis Obispo County coastal area, 238
Costs: of planting two domestic trout strains at Lake Berryessa, 44-49
Coyote: food habits in Sagehen Creek basin, 4-12
Coyote, juvenile: food habits at Sagehen Creek basin, 9-11
Crab, box: case of symbiosis with blacktail snailfish, 239-240
Crab, umbrella: range extension, 240-243
Croaker, white: DDT residues, 22-26, 198-203
Cycling, ovarian: of gobies at the Salton Sea, 50-57
DDT: in the diet of pheasants producing melanistic mutants, 175-178; residues in California marine fishes, 22-26; residues in white croakers, 198-203; toxicity to surfperch, 127-132
Deer: as food for coyotes, 6-7; occurrence of rabies, 141-144; study of Mineral King herd, 133-140
Dieldrin: toxicity to surfperch, 127-132
Disease: occurrence of rabies in deer, 141-144
Distribution, geographic: appearance of California scorpionfish in Monterey Bay, 238-239; extension of range of yellowfin goby, 326-327; of flag and redbanded rockfishes, 32-36; of the cottonmouth jack, 245-246; of the loggerhead, 248; of the umbrella crab, 240-243; of white and green sturgeons, 102-106; range extension for four surfperches, 27-31; range ex-

- tension of *Palaemonetes paludosus*, 154-155
- Distribution, vertical: of euphausiids off the coast of Washington, 179-190
- Eggs, fish: effect of salinity on Salton Sea fish, 58-66
- Elkhorn Slough: yellowfin goby occurrence, 326-327
- Embiotocidae: geographical records for four species, 27-31
- Endrin: toxicity to surfperch, 127-132
- Estuary, Sacramento-San Joaquin: white sturgeon population study, 94-101; migration study of white and green sturgeons, 102-106
- Euphausiids: diel changes in the vertical distribution, 179-190
- Fauna, stream: effects of diesel fuel, 291-295
- Feeding: of spiny lobster phyllosomes in the laboratory, 107-115
- Fishery: characteristics at Folsom Lake, 75-93; for northern anchovy during 1969-70 season, 121-126; for trout at Lake Berryessa, 44-49; for white sturgeon, 99-100; of Pacific mackerel in southern California 1964-65 through 1967-68 seasons, 13-21; of Pacific mackerel 1968-69 and 1969-1970 seasons, 116-120
- Fishes, intertidal: of Trinidad Bay and vicinity, 315-320
- Folsom Lake: angling quality study, 75-93; site of primary productivity studies, 254-267
- Food habits: of coyotes in Sagehen Creek basin, 4-12; of Klamath River fall-run steelhead, 217-218; of San Joaquin kit fox, 167-169; of trout in high elevation Sierra lakes in winter, 231-237
- Fox, San Joaquin kit: life history, 162-174
- Fuel, diesel: effects of spilling into stream, 291-295
- Goby, longjaw: ovarian cycling at the Salton Sea, 50-57
- Goby, yellowfin: southern range extension, 326-327
- Growth: characteristics of Klamath River steelhead, 207-211; of spiny lobster phyllosomes in laboratory, 107-115; of Piute sculpin, 287
- Half-pounder: steelhead of the Klamath River, 204-220
- Harvest: of two domestic trout strains at Lake Berryessa, 44-49
- Insecticides, organochlorine: toxicity to surfperch, 127-132
- Insects: as food for coyotes, 9
- Invertebrates, aquatic: capture by use of polypropylene light trap, 149-152
- Jack, cottonmouth: added to marine fauna of California, 245-246
- Klamath River: characteristics of fall-run steelhead, 204-220
- Lake Berryessa: study of planting two domestic strains of rainbow trout, 44-49
- Larvae, fish: effect of salinity on Salton Sea fish, 58-66
- Length: composition of anchovies taken 1969-70 season, 121-126
- Life history: of Piute sculpin, 285-290; of San Joaquin kit fox, 162-174
- Lingcod: DDT residues, 22-26
- Lions, sea: predation on salmon, 37-43
- Lobster, California spiny: molting and growth of phyllosomes in laboratory, 107-115
- Logperch: range extension, 248
- Mackerel, jack: DDT residues, 22-26
- Mackerel, Pacific: age composition of the catch 1964-65 through 1967-68 seasons, 13-21; southern California fishery during 1968-69 and 1969-70 seasons, 116-120
- Merle Collins Reservoir: bass tagging study, 145-147; site of largemouth bass mortality and survival study, 221-230; site of primary productivity studies, 254-267
- Methods: aging Pacific mackerel 1964-65 through 1967-68 seasons, 13-21; constructing polypropylene light trap for aquatic invertebrates, 149-152; coyote food habits study, 4-12; determining DDT residues in marine fishes, 22-26; determining DDT residues in white croakers, 199; determining effects of salinity on eggs and larvae of Salton Sea fish, 58-66; determining organochlorine toxicity to surfperch, 127-132; ovarian cycling of longjaw goby, 50-57; rearing spiny lobster phyllosomes in laboratory, 107-115; separation of *Sebastes babcocki* and *S. rubrivinctus*, 32-36; studying life history of the kit fox, 162-174; studying mortality and survival of bass at Merle Collins Reservoir, 221-230 studying vertical distribution of euphausiids, 179-190; study of Mineral King deer herd, 133-140; study of fall-run steelhead of the Klamath River, 204-220; study of life history of Piute sculpin, 285-290; study of Modoc sucker, 277-284; study of predation by sea lions on salmon, 37-43; study of subpopulations of Pacific sardines, 296-314; study of reservoir primary productivity, 254-267; study of trout fishery at Lake Berryessa, 44-49; study of variation in the swell shark, 191-197; tagging sturgeon to provide migration data, 102-106; tagging to determine white sturgeon population,

- 94-101; testing tolerance of pileworm to salinity, 152-154; use of roving creel census at Folsom Lake, 75-93
- Migration: of Klamath River steelhead, 211-214; of Mineral King deer, 133-140; of white and green sturgeons, 102-106
- Mineral King: deer study, 133-140
- Molting: of spiny lobster phyllosomes in laboratory, 107-115
- Monterey Bay: study of predation by sea lions on salmon, 37-43
- Morphology: of the Modoc sucker, 277-284
- Mortality rates: of largemouth bass at Merle Collins Reservoir, 221-230
- Mt. Whitney trout strain: performance at Lake Berryessa, 44-49
- Mullet, striped: apparent spawning at sea, 246-248
- Mutant: melanism in ringneck pheasants, 175-178
- Net: midwater trawl for taking threadfin shad, 268-276
- Perch, dwarf: pesticide toxicity studies, 127-132
- Perch, shiner: pesticide toxicity studies, 127-132; southern range record, 28
- Pesticides: chlorinated hydrocarbon residues in California marine fishes, 22-26; toxicity of organochlorines to surfperch, 127-132
- Pheasant, ringneck: appearance of melanistic mutant, 175-178
- Phyllosomes: growth of spiny lobsters under laboratory conditions, 107-115
- Pileworm: tolerance of high salinity, 152-154
- Pollution: effects of diesel fuel on a stream fauna, 291-295
- Population: of Mineral King deer herd, 133-140; resurgence in sharpnose seaperch, 28-31; study of variation within subpopulations of Pacific sardines, 296-314; survey of condors, 1971, 327-328; white sturgeon in Sacramento-San Joaquin estuary, 94-101
- Predation: by sea lions on salmon, 37-43
- Productivity, primary: in a new and old California reservoir, 254-267
- Rabbits: as food for coyotes, 8
- Rabies: occurrence in deer, 141-144
- Raptors: numbers seen during 1970 condor survey, 67-68; seen during 1971 condor survey, 327-328
- Rattail, roughscale: DDT residues, 22-26
- Reproduction: failure of pelagic cormorants to breed, 238; of longjaw goby at the Salton Sea, 50-57; of Piute sculpin, 287
- Reviews: A trout and salmon fisherman for seventy-five years, 330; Biology and water pollution control, 249; British Columbia game fish, 71; Come wade the river, 71; Fishes of Montana, 329-330; Fishless days, angling nights, 250; Hardy's book of fishing, 331; Hikers and back-packers handbook, 70; If deer are to survive, 157; Kamloops, 71; Life and death in a coral sea, 250; Mountain sheep: a study in behavior and evolution, 329; Remembrances of rivers past, 330; Round river, 250; Sea shells of tropical west America; marine molluscs from Baja California to Peru, 249; Systematics, variation, distribution, and biology of rockfishes of the subgenus *Sebastomus* (Pisces, Scorpaenidae, *Sebastes*), 156-157; The art and science of fly fishing, 321; The dry fly and fast water and the salmon and the dry fly, 330-331; The ecology of running waters, 156; The New York aquarium book of the water world, 332; The vanishing jungle—the story of the World Wildlife Fund expeditions to Pakistan, 70-71; Wildlife of Mexico: the game birds and mammals, 250; World dynamics, 332
- Rockfish, flag: identity and range, 32-36
- Rockfish, redbanded: identity and range, 32-36
- Rodents: as food for coyotes, 8
- Rusk Creek: site of study of Modoc suckers, 277-284
- Sablefish: DDT residues, 22-26
- Sagehen Creek: site of coyote food habits study, 4-12; study of life history of Piute sculpin, 285-290
- Salinity: tolerance by pileworm, 152-154
- Salmon: study of predation by sea lions, 37-43
- Salton Sea: ovarian cycling of longjaw goby, 50-57; study of salinity stress on fish eggs and larvae, 58-66; tests on pileworm tolerance of salinity, 152-154
- San Clemente Island: survival of Guadalupe cardinalfish, 68-69
- Sanddab, Pacific: DDT residues, 22-26
- Sardine, Pacific: subpopulation study, 296-314
- Sargo: effect of salinity on eggs and larvae, 58-66
- Scorpionfish, California: reoccurrence in Monterey Bay, 238-239
- Sculpin, Piute: life history, 285-290
- Seaperch, pink: host for parasitic copepod, 323-325
- Seaperch, rubberlip: southern range record, 28

- Seaperch, sharpnose: southern range records and notes on population resurgence, 28-31
- Sea urchin: in association with acorn barnacles, 321-323
- Shad, threadfin: trawl designed for catching, 268-276
- Shark, swell: population differences off California and Baja California, 191-197
- Sheep: as food for coyotes, 7
- Snailfish blacktail: case of symbiosis with box crab, 239-240
- Sole, butter: case of reversal, 244, 245
- Sole, Dover: tailless specimens taken off Oregon, 147-148
- Sole, English: DDT residues, 22-26
- Sole, petrale: DDT residues, 22-26
- Spawning: of striped mullet at sea, 246-248
- Stress, salinity: on Salton Sea fish eggs and larvae, 58-66
- Sunfish: fishery at Folsom Lake, 84
- Sturgeon, green: migration, 102-106
- Sturgeon, white: migration, 102-106; population in Sacramento-San Joaquin estuary, 94-101
- Sucker, Modoc: morphology, variation, feeding adaptations, 277-284
- Surfperch: pesticide toxicity studies, 127-132
- Surfperch, silver: southern range records, 27
- Surfperch, spotfin: host for parasitic copepod, 323-325
- Survey: 1970 California condor population, 67-68
- Survival: of Guadalupe cardinalfish at San Clemente Island, 68-69; of two domestic trout strains at Lake Berryessa, 44-49
- Survival rates: of largemouth bass at Merle Collins Reservoir, 221-230
- Symbiosis: between blacktail snailfish and box crab, 239-240
- Tagging: of largemouth bass at Merle Collins Reservoir, 145-147; to determine sturgeon migration, 102-106; to determine white sturgeon populations, 94-101
- Telemetry: use in studying kit foxes, 165; use in studying Mineral King deer, 138-140
- Toxicity: of organochlorines to surfperch, 127-132
- Trap: for aquatic invertebrates, 140-152
- Trawl, midwater: for threadfin shad, design and operation, 268-276
- Trinidad Bay: intertidal fishes, 315-320
- Trout, rainbow: fishery at Folsom Lake, 85; performance of two domestic strains at Lake Berryessa, 44-49
- Trout, steelhead: characteristics of fall-run in Klamath River, 204-220
- Trout: winter food habits in high elevation Sierra lakes, 231-237
- Vegetable matter: in coyote food habits, 9, 11

O